

Nonchemical methods for rice crop insect pest management in Africa

Silvie¹, Pierre, Adda², Cyrille, Togola², Abou, Nwilene³, Francis, Menozzi⁴, Philippe

¹CIRAD UPR Systèmes de culture annuels, F-34398 Montpellier Cedex 5, France, pierre.silvie@cirad.fr

²Africa Rice Center (AfricaRice), 01 B.P. 2031, Cotonou, Benin, West Africa

³Africa Rice Center (AfricaRice), Oyo Road, PMB 5320, Ibadan, Nigeria

⁴CIRAD, UPR SCA, F-34398 Montpellier Cedex 05, France ; Africa Rice Center, 01 BP 2031, Cotonou, Benin

Keywords

Pest management, organic rice, Africa

Introduction

Various rice growing methods are implemented in sub-Saharan Africa. The type of method depends mainly on the irrigation possibilities: rainfed upland and lowland rice, irrigated rice. In the field, amongst the different factors that hamper crop growth and production, insects are often ranked after other constraints associated with water management, low soil fertility, weed infestation (at emergence), nematodes, diseases and birds (at harvest) (AfricaRice, 2011). During storage, Curculionidae species such as *Sitophilus oryzae* are known to be problematic, but an increase in seed production could also give rise to serious problems associated with lepidopterans such as *Sitotroga cerealella* (Togola et al., 2010). In Africa, the identity and diversity of rice pests and their natural enemies in Africa are well known at the field level (Brenière, 1982).

We conducted a survey to determine what nonchemical methods are currently available for managing rice insect pests in sub-Saharan African rice-producing countries (excluding Madagascar).

What has been published about organic rice in Africa?

A search in databases, including the Web of Science (online since 1975) and SCOPUS (since 1960), and the OvidSP platform (queries of AGRICOLA/1970, AGRIS/1991, CAB/1973, ECONLIT/1960 and PASCAL/1984), was carried out to highlight bibliographical references concerning organic rice. This search, using the expression 'ORGANIC RICE', was done on 27 March 2012. A boolean query with the added words 'AFRICA' or 'MADAGASCAR' was also conducted. Two hundred and three references were found when duplicate references were eliminated. Thirty of these concerned insects, mainly in Asia. No literature references on organic rice in Africa were found. The information gathered and presented in this short summary is thus derived from articles published by research experts on IPM in rice cropping conditions in Africa. Only insects causing crop damage in the field were taken into consideration, not those that infest stored rice.

Results on nonchemical pest management methods

Nonchemical management methods that were studied:

- The use of partially resistant varieties. There are indeed between-variety differences in the susceptibility to some pests such as termites (Agunbiade et al., 2009), the Cecidomyidae fly *Orseolia oryzivora*, which is often considered to be a serious pest (Williams et al., 2002; Nwilene et al., 2002) and stemborers (Nwilene et al., 2011). Tolerant varieties do not hamper insect development. Little is known about tolerance and resistance mechanisms except the work by Nwilene et al. (2009). The overcompensation capacities of some varieties, i.e. greater tillering, are involved.
- Strip-cropping rice with maize. Planting rows of rice between four rows of maize (all rows of equal width) leads to a reduction in attacks of stemborers such as *Maliarpha separata* and

Sesamia calamistis on the rice crops (Nwilene et al., 2011). Diversion of *Sesamia* stemborers towards the maize crop is the suggested mechanism.

- The application of plant extracts such as neem (*Azadirachta indica*) oil has been studied and recommended for controlling termites (Nwilene et al., 2008a).
- Regular weeding and the use of the entomopathogenic fungus *Metarhizium anisopliae* result in lower termite population and damage in rice field (Togola et al., 2012).
- The management of habitats in the immediate vicinity of plots could be carried out by two different approaches: (i) preserving the diversity of natural enemies, predators or parasitoids: planting *Paspalum scrobiculatum* (Poaceae) along the edges of rice crop fields enables the development of *Orseolia bonzii* (Cecidomyidae), which does not harm rice crops. This species is a substitute host of *O. orseolia* parasitoids such as *Platygaster diplosisae* and *Aprostocetus procerae* (Nwilene et al., 2008b); and (ii) eliminating host plants or rice ratoons that facilitate the development or survival of some pests like Diopsides flies (Togola et al., 2011) or gall midges (Williams et al., 2002).

Habitat management on a broader landscape scale has not been studied in Africa. This is of considerable research interest, in line with the studies carried out in Japan by Takada et al. (2012) on the bug *Stenotus rubrovittatus* (Miridae), pests of seeds.

Hence, if certain rules are actually fulfilled, e.g. with respect to organic fertilization or crop rotations, several techniques are potentially available for organic rice production in Africa.

References

- [1] Africa Rice Center (AfricaRice), 2011. Boosting Africa's rice sector: A research for development strategy 2011-2020. Cotonou, Benin.
- [2] Agunbiade, TA, et al., 2009. Resistance status of Upland NERICA rice varieties to termite damage in Northcentral Nigeria. *Journal of Applied Science* 9, 3864-3869.
- [3] Brenière, J, 1982. Estimation des pertes dues aux ravageurs du riz en Afrique de l'Ouest. *Entomophaga* 27 (H.S.), 71-80.
- [4] Nwilene, F, et al., 2002. Reactions of differential rice genotypes to African rice gall midge in West Africa. *International Journal of Pest Management* 48, 195-201.
- [5] Nwilene, F, et al., 2008a. Efficacy of traditional practices and botanicals for the control of termites on rice at Ikenne, southwestern Nigeria. *International Journal of Tropical Insect Science* 28, 37-44.
- [6] Nwilene, F, et al., 2008b. Parasitoid biodiversity conservation for sustainable management of the African rice gall midge, *Orseolia oryzivora* (Diptera: Cecidomyidae) in lowland rice. *Biocontrol Science and technology* 18, 1075-1081.
- [7] Nwilene, F, et al., 2009. Antixenosis component of rice resistance to African rice gall midge, *Orseolia oryzivora*. *International Rice Research Notes* 34, 1-5.
- [8] Nwilene, F, et al., 2011. Effect of intercropping maize and cassava with Upland NERICA rice varieties on stemborer attacks in Southwestern Nigeria. *Journal of entomology* 8, 417-48.
- [9] Takada, MB, et al., 2012. Multiple spatial scale factors affecting mired bug abundance and damage level in organic rice paddies. *Biological Control* 60, 169-174.
- [10] Togola, A, et al., 2010. Présence, populations et dégâts de l'alucite des céréales *Sitotroga cerealella* (Olivier) (Lepidoptera, Gelechiidae) sur les stocks de riz au Bénin. *Cahiers Agricultures* 19: 205-209. doi: 10.1684/agr.2010.0398.
- [11] Togola, A, et al., 2011. Sédentarisation des populations des mouches Diopsides dans les agro-systèmes rizicoles au Bénin. *Tropicicultura* 29, 101-106.
- [12] Togola, A, et al., 2012. Effect of upland rice varieties and cultural practices on termite populations and damage in the field. *Journal of Applied Sciences* (in press). doi: 10.3923/jas.2012.
- [13] Williams CT, et al., 2002. African rice gall midge research guide. Bouaké, Côte d'Ivoire: West Africa Rice development Association, and Wallingford, UK: CAB International, 28 p.

Appendix Table 1 The main rice pests and nonchemical management methods

Physiological stages (from Brenière, 1982)	Families and species	Nonchemical pest control methods and efficacy ^(a)	References
Herbaceous state (nursery or 1 st month)	Coccinellidae (<i>Chnootriba similis assimilis</i> , <i>Epilachna reticulata</i>) Pyralidae (<i>Nymphula depunctalis</i>)		
Root pests	Termites	Varietal tolerance (+) Neem seed oil (++) Regular weeding (++) <i>Metarhizium anisopliae</i> (++)	[2] [5] [12] [12]
	Scarabaeidae (<i>Heteronychus</i> spp.) Nematodes		
<u>Tillering</u> Foliage	Arctiidae (<i>Diacrisia scortilla</i>) Hesperiidae (<i>Parnara</i> sp., <i>Borbo</i> sp.) Noctuidae (<i>Spodoptera</i> spp.) Pyralidae <i>Marasmia trapezalis</i> Hispidae (<i>Hispa unsambarica</i> , <i>Lema</i> spp., <i>Trichispa sericea</i>)	Varietal tolerance (+)	
Sheaths and stems	Cecidomyiidae (<i>Orseolia oryzivora</i>) Diopsidae (<i>Diopsis apicalis</i> , <i>D. thoracica</i>) Crambidae (<i>Chilo zacconius</i>) Pyralidae (<i>Eldana saccharina</i> , <i>Maliarpha separatella</i>)	Varietal tolerance (++) Parasitoid conservation (++) Ratoon elimination (++) Strip-cropping (maize) (++)	[4,7] [6] [11] [8]
<u>Heading</u> Stem-borers	Crambidae (<i>Chilo zacconius</i>) Noctuidae (<i>Sesamia calamistis</i> , <i>S. nonagrioides botanephaga</i> , <i>S. nonagrioides penniseti</i> , <i>S. poephaga</i> , Pyralidae (<i>M. separatella</i> , <i>Scirpophaga</i> spp.)	Varietal tolerance (+) Strip-cropping (maize)(++)	[4] [8]
<u>Ripening</u> Stem-borers	Noctuidae (<i>Sesamia</i> spp.) Pyralidae (<i>Maliarpha separatella</i> , <i>Scirpophaga</i> spp.) Pentatomidae, Lygaeidae	Varietal tolerance (+)	[4]
Sucking insects	Delphacidae, Jassidae	Varietal tolerance (+++)	

^(a) +, ++, +++: low to high efficacy